AD-A197 466

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION: Unclassified	1b RESTRICTIVE MARKINGS:
2a SECURITY CLASSIFICATION AUTHORITY	3 DISTRIBUTION / AVAILABILITY OF REPORT: Approved for public release; distribution unlimited.
2b DECLASSIFICATION / DOWNGRADING SCHEDULE	
4 PERFORMING ORGANIZATION REPORT NUMBER(S): Technical Report No. DU/DC/TR-08	5 MONITORING ORGANIZATION REPORT NUMBER(S):
6a NAME OF PERFORMING ORGANIZATION: Department of Chemistry, Duke University; 6b OFFICE SYMBOL (If applicable):	7a NAME OF MONITORING ORGANIZATION: Office of Naval Research
6c ADDRESS (City, State, and ZIP Code): Durham, NC 27706	7b ADDRESS (City, State, and ZIP Code): 800 N. Quincy Street, Arlington, VA 22217-5000
8a NAME OF FUNDING / SPONSORING ORGANIZATION: Office of Naval Research; 8b OFFICE SYMBOL (If applicable):	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER: N00014-83-K-0572 R&T Code 4135008---02
8c ADDRESS (City, State, and ZIP Code): 800 N. Quincy Street, Arlington, VA 22217-5000	10 SOURCE OF FUNDING NUMBERS: PROGRAM ELEMENT NO: NR; PROJECT NO: 053; TASK NO: 841; WORK UNIT ACCESSION NO:

11 TITLE (Include Security Classification)
The Twist-Boat Conformation in a Gallium-Arsenic Six-Membered Ring: Crystal Structure of $[(Me_3SiCH_2)_2AsGaBr_2]_3$

12 PERSONAL AUTHOR(S)
R.L. Wells, A.P. Purdy, A.T. McPhail, and C.G. Pitt

13a TYPE OF REPORT	13b TIME COVERED	14 DATE OF REPORT (Year, Month, Day)	15 PAGE COUNT
Technical	FROM _____ TO _____	88/07/05	12

16 SUPPLEMENTARY NOTATION
Accepted for Publication in Journal of Organometallic Chemistry

17 COSATI CODES		
FIELD	GROUP	SUB-GROUP

18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)
Organogallium-arsenic, Crystal Structure, Conformation, Gallium-arsenic six-membered ring.

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

$[(Me_3SiCH_2)_2AsGaBr_2]_3$, the first trimeric mono(arsino)gallane to be structurally characterized by X-ray crystallographic analysis, has been shown to contain a $(Ga-As)_3$ ring in the twist-boat conformation. Crystals of the trimer are monoclinic, space group $P2/c(C_{2h}^4)$, with a 18.877(4), b 13.710(3), c 24.821(4) Å, β 122.24(2)°, U 5433.4 Å^3, and Z 4. The Ga-As bond lengths [2.432(2)-2.464(1) Å] are all less than any of those found in the $(Ga-As)_2$ rings of dimeric arsinogallanes.

20 DISTRIBUTION / AVAILABILITY OF ABSTRACT	21 ABSTRACT SECURITY CLASSIFICATION
☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS	Unclassified
22a NAME OF RESPONSIBLE INDIVIDUAL: Richard L. Wells	22b TELEPHONE (Include Area Code): (919) 684-6404 / 22c OFFICE SYMBOL:

DD FORM 1473, 84 MAR — 83 APR edition may be used until exhausted. All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE: Unclassified

DTIC ELECTE JUL 08 1988

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0572

R&T Code 4135008---02

Technical Report No. DU/DC/TR-08

The Twist-Boat Conformation in a Gallium-Arsenic Six-Membered Ring: Crystal Structure of $[(Me_3SiCH_2)_2AsGaBr_2]_3$

by

R. L. Wells, A. P. Purdy
A. T. McPhail, and C. G. Pitt

Prepared for Publication in
Journal of Organometallic Chemistry

Duke University
Department of Chemistry
Durham, NC 27706

Accession For
NTIS GRA&I [X]
DTIC TAB [ ]
Unannounced [ ]
Justification
By
Distribution/
Availability Codes
Dist | Avail and/or Special
A-1

July 5, 1988

DTIC COPY INSPECTED 6

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited

THE TWIST-BOAT CONFORMATION IN A GALLIUM-ARSENIC SIX-MEMBERED RING: CRYSTAL STRUCTURE OF $[(Me_3SiCH_2)_2AsGaBr_2]_3$

RICHARD L. WELLS*, ANDREW P. PURDY, ANDREW T. McPHAIL, and COLIN G. PITT*

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University, Durham, NC 27706 (U.S.A.)

SUMMARY

$[(Me_3SiCH_2)_2AsGaBr_2]_3$, the first trimeric mono(arsino)gallane to be structurally characterized by X-ray crystallographic analysis, has been shown to contain a $(Ga-As)_3$ ring in the twist-boat conformation. Crystals of the trimer are monoclinic, space group $P2/c(C_{2h}^4)$, with a 18.877(4), b 13.710((3), c 24.821(4) Å, β 122.24(2)°, U 5433.4 Å^3, and Z 4. The Ga-As bond lengths [2.432(2)-2.464(1) Å] are all less than any of those found in the $(Ga-As)_2$ rings of dimeric arsinogallanes.

Introduction

The facile preparation of a number of new gallium-arsenic compounds, including $[(Me_3SiCH_2)_2AsGaBr_2]_3$ (1), using silylarsines was recently demonstrated by work in our laboratories [1], and 1 is one of a group of mono(arsino)gallanes now known to exist as trimers on the basis of NMR and cryoscopic molecular weight data. Some in this group are reported to be solely trimeric in solution, e.g., $(Me_2AsGaMe_2)_3$ [2], $[(Me_3SiCH_2)_2AsGaCl_2]_3$

[1a], and 1 [1b], whereas dimer-trimer equilibria have been shown to be present in solutions of others, e.g., $[(Ph)Me_3SiCH_2AsGaMe_2]_{2,3}$ [1d], $[(Ph)Me_3SiCH_2AsGaPh_2]_{2,3}$ [1d], $[(Me_3SiCH_2)_2AsGaMe_2]_{2,3}$ [1d], $[(Me_3SiCH_2)_2AsGaMe(Cl)]_{2,3}$ [1b], and $[(Me_3SiCH_2)_2AsGaPh(Cl)]_{2,3}$ [1b]. Notwithstanding the preceding information, no structural details for a trimeric mono(arsino)gallane have appeared in the literature to date. Thus, a single-crystal X-ray analysis was performed on 1 to determine the geometrical features of the six-membered $(Ga-As)_3$ ring in this type of molecule.

Experimental

Preparation of $[(Me_3SiCH_2)_2AsGaBr_2]_3$ (1) [1b]

Compound 1 was prepared from the reaction of $(Me_3SiCH_2)_2AsSiMe_3$ (0.20 g, 0.65 mmol) with $GaBr_3$ (0.21 g, 0.65 mmol) in ligroin (~20 mL, bp 95-110°), rather than in pentane which was used in the initially reported synthesis [1b], and it precipitated from the solution as small, colorless crystals in contrast to the large crystals which separated from pentane [1b]. Removal of the mother liquor followed by drying under vacuum resulted in a large number of small, uniformly sized crystals suitable for an X-ray study. The ^1H NMR spectrum of a sample of the latter crystals in C_6D_6 was identical to that of a sample of the large crystals [1b].

Crystal data. $C_{24}H_{66}As_3Br_6Ga_3Si_6$, M 1436.69, monoclinic, space group $P2/c(C_{2h}^4)$, a 18.877(4), b 13.710(3), c 24.821(4) Å, β 122.24(2)°, U 5433.4 Å^3, Z 4, D_c 1.756 g cm^{-1}, μ(Cu-Kα radiation, λ 1.5418 Å) 104.6 cm^{-1}, sample dimensions 0.15 x 0.25 x 0.54 mm (sealed inside a thin-walled capillary).

Crystallographic measurements. Preliminary unit-cell parameters and space

group information were obtained from oscillation and Weissenberg photographs. Intensity data ($+h$, $+k$, $\pm l$, 9668 independent reflections) were recorded on an Enraf-Nonius CAD-4 diffractometer(Cu-$K\alpha$ radiation, incident-beam monochromator; ω-2θ scans, $\theta_{max.} = 67^{\circ}$). Those 4798 reflections with $I > 3.0\sigma(I)$ were retained for the structure analysis and corrected for the usual Lorentz and polarization effects. An empirical absorption correction, based on the ϕ-dependence of the intensities of 4 reflections with χ ca. 90°, was also applied to the data. Refined unit-cell parameters were derived from the diffractometer setting angles for 25 reflections ($40^{\circ} < \theta < 55^{\circ}$) widely separated in reciprocal space.

Structural analysis. The structure was solved by direct methods. Full-matrix least-squares adjustment of non-hydrogen atom positional and anisotropic thermal parameters, with hydrogen atoms included at their calculated positions in the later iterations, converged to $R = 0.045$ ($R_w = 0.060$) {$R = \Sigma||F_o| - |F_c||/\Sigma|F_o|$; $R_w = [\Sigma w(|F_o| - |F_c|)^2/\Sigma w|F_o|^2]^{1/2}$}. For the structure-factor calculations, neutral atom scattering factors and their anomalous dispersion corrections were taken from ref. 3. In the least-squares iterations, $\Sigma w\Delta^2$ [$w = 1/\sigma^2(|F_o|)$, $\Delta = (|F_o| - |F_c|)$] was minimized. All crystallographic calculations were performed on a PDP11/44 computer by use of the Enraf-Nonius SDP suite of programs incorporating the direct methods program MULTAN11/82. Final non-hydrogen atom positional parameters are in Table 1 and selected bond lengths and angles are in Table 2. Tables of interatomic distances and angles, torsion angles, thermal parameters, hydrogen atom parameters, and observed and calculated structure factors are available from the authors (R.L.W.).

Discussion

Crystals of 1 contain two crystallographically independent, but structurally identical, discrete molecules having the constitution illustrated in Figure 1. Each of the molecules lies across a crystallographic $\underline{C}_2$ axis of symmetry passing through As(1) and Ga(4) and, with the endocyclic torsion angles also related by an approximate $\underline{C}_2$ axis passing through the mid-points of the Ga(2)-As(3) and Ga(2')-As(3') bonds, the $(Ga-As)_3$ ring adopts the somewhat rare twist-boat conformation (Figure 2). The torsion angles of -30.2, +59.9, and -25.0° † [Ga(2')-As(1)-Ga(2)-As(3), As(1)-Ga(2)-As(3)-Ga(4), and Ga(2)-As(3)-Ga(4)-As(3'), respectively] compare favorably to the theoretical values of -31, +65, and -31° for equivalent torsion angles in the twist-boat form of cyclohexane [4]. In $(\underline{t}\text{-}Bu_2SbGaCl_2)_3$, the $(Ga-Sb)_3$ ring has a somewhat irregular boat conformation, whereas in $(Me_2AsInMe_2)_3$, the $(In-As)_3$ ring is found to be both planar and puckered [5].

Endocyclic bond angles at As [102.88(7), 110.47(5)°] differ significantly and both are smaller than those subtended at Ga [113.46(8)], 121.36(7)°] which are also unequal. Exocyclic Br-Ga-Br angles [112.98(8), 111.60(9)°] differ only slightly and the C-As-C angles [108.5(4), 108.4(5)°] are equal. Ga-As bond lengths [2.432(2) - 2.464(1) Å] are all less than any of the distances found thus far between four-coordinate Ga and four-coordinate As centers in dimeric organogallium compounds [1b, 6-8] for which the shortest

---

† Averages for the two crystallographically-independent molecules are cited; similarly, all other numbers in the discussion section pertaining to 1 are average values. Where different, the higher of two estimated standard deviations are quoted.

[2.513(1)°] occurs in the planar $(Ga-As)_2$ ring of $\{[(Me_3SiCH_2)_2As]_2GaBr\}_2$ [1b] and the longest [2.581(1) Å] is present in $\{[(Me_3SiCH_2)_2As]_3Ga\}_2$ which contains a nonplanar $(Ga-As)_2$ ring [7]. Interestingly, the shortest Ga-As distance in 1 [2.432(2) Å] is not significantly different from the shortest previously reported length for an organogallium-arsenic compound, _viz._, the Ga-_exo_-As distance [2.437(1) Å] between four-coordinate Ga and three-coordinate As in the dimer $\{[(Me_3SiCH_2)_2As]_2GaBr\}_2$ [1b].

Acknowledgement.

We thank the Office of Naval Research and the Duke University Research Council for financial support.

References

1 (a) C. G. Pitt, A. P. Purdy, K. T. Higa, and R. L. Wells, Organometallics, 5 (1986) 1266. (b) A. P. Purdy, R. L. Wells, A. T. McPhail, and C. G. Pitt, Organometallics, 6 (1987) 2099. (c) R. L. Wells, S. Shafieezad, A. T. McPhail, and C. G. Pitt, J. Chem. Soc., Chem. Commun., (1987) 1823. (d) A. P. Purdy, Ph. D. Dissertation, Duke University, Durham, NC 1987. (e) R. L. Wells, W. K. Holley, S. Shafieezad, A. T. McPhail, and C. G. Pitt, Fifth International Symposium on Inorganic Ring Systems, Amherst, MA, (1988); Phosphorus and Sulfur (to be published).

2 O. T. Beachley and G. E. Coates, J. Chem. Soc., (1965) 3241.

3 International Tables for X-ray Crystallography, Vol IV, The Kynoch Press, Birmingham, England, (1974).

4 R. Bucourt, Topics in Stereochemistry, 8 (1974) 159 (E. L. Eliel and N. L. Allinger, Editors), Interscience, New York, NY.

5 A. H. Cowley, R. A. Jones, K. B. Kidd, C. M. Nunn, and D. L. Westmoreland, J. Organomet. Chem., 341 (1988) C1.

6 R. L. Wells, A. P. Purdy, A. T. McPhail, and C. G. Pitt, J. Organomet. Chem., 308 (1986) 281.

7 R. L. Wells, A. P. Purdy, K. T. Higa, A. T. McPhail, and C. G. Pitt, J. Organmet. Chem., 325 (1987) C7.

8 A. M. Arif, B. L. Benac, A. H. Cowley, R. Geerts, R. A. Jones, K. B. Kidd, J. M. Power, and S. T. Schwab, J. Chem. Soc., Chem. Commun., (1986) 1543.

TABLE 1. NON-HYDROGEN ATOM FRACTIONAL COORDINATES ($X10^4$), WITH ESTIMATED STANDARD DEVIATIONS IN PARENTHESES

Molecule I

Atom	x	y	z
As(1)[a]	0(-)	2200.9(11)	2500(-)
Ga(2)	-0.6(6)	3301.9(9)	1732.2(5)
As(3)	-658.5(6)	4919.9(8)	1521.7(4)
Ga(4)[a]	0(-)	5905.8(13)	2500(-)
Br(21)	-390.8(7)	2500.8(10)	789.6(5)
Br(22)	1393.4(6)	3837.4(10)	2286.2(6)
Br(41)	-1044.1(7)	6861.4(10)	2456.6(7)
Si(11)	-1229(2)	435(2)	2548(1)
Si(31)	-580(2)	6668(3)	571(2)
Si(32)	-2655(2)	4241(2)	482(1)
C(111)	-977(5)	1381(7)	2110(4)
C(112)	-1785(6)	1066(10)	2885(5)
C(113)	-1910(7)	-458(9)	1927(6)
C(114)	-283(7)	-181(10)	3211(6)
C(311)	-334(7)	5405(8)	950(5)
C(312)	431(8)	7219(10)	773(6)
C(313)	-1092(8)	7489(9)	877(6)
C(314)	-1261(10)	6489(12)	-308(6)
C(321)	-1869(5)	5073(8)	1109(5)
C(322)	-2533(7)	2996(9)	806(6)
C(323)	-3689(7)	4739(9)	260(6)
C(324)	-2570(7)	4235(10)	-224(5)

Molecule II

Atom	x	y	z
As(1)[a]	5000(-)	-2364.5(10)	2500(-)
Ga(2)	4913.7(6)	-1254.7(9)	1702.7(5)
As(3)	4259.6(5)	357.9(8)	1517.6(4)

TABLE 1. (CONTINUED)

Atom	x	y	z
Ga(4)[a]	5000(-)	1343.6(12)	2500(-)
Br(21)	4486.8(8)	-2032.0(10)	746.4(5)
Br(22)	6302.4(6)	-707.8(10)	2224.1(6)
Br(41)	3900.2(7)	2292.4(9)	2515.1(6)
Si(11)	3820(2)	-4112(3)	2628(2)
Si(31)	4254(2)	2104(3)	540(1)
Si(32)	2257(2)	-341(2)	523(1)
C(111)	4016(5)	-3194(8)	2157(5)
C(112)	3362(8)	-3503(12)	3045(6)
C(113)	3059(8)	-4975(11)	2010(7)
C(114)	4768(8)	-4791(10)	3214(6)
C(311)	4508(6)	845(8)	895(4)
C(312)	5227(7)	2672(10)	688(6)
C(313)	3763(8)	2916(9)	867(6)
C(314)	3525(10)	1955(11)	-345(7)
C(321)	3075(5)	499(8)	1162(4)
C(322)	2407(7)	-1580(8)	854(6)
C(323)	1225(6)	121(9)	316(5)
C(324)	2317(7)	-347(9)	-201(5)

[a]Occupancy factor = 0.50

TABLE 2. SELECTED BOND LENGTHS (Å) AND ANGLES (deg.), WITH ESTIMATED STANDARD DEVIATIONS IN PARENTHESES

	Molecule I	Molecule II
(a) Bond Lengths		
As(1)-Ga(2)	2.431(2)	2.432(2)
As(1)-C(111)	1.923(9)	1.945(9)
Ga(2)-As(3)	2.458(2)	2.453(2)
Ga(2)-Br(21)	2.320(2)	2.319(2)
Ga(2)-Br(22)	2.344(1)	2.343(1)
As(3)-Ga(4)	2.459(1)	2.468(1)
As(3)-C(311)	1.939(14)	1.956(13)
As(3)-C(321)	1.954(9)	1.926(9)
Ga(4)-Br(41)	2.321(2)	2.325(2)
(b) Bond Angles		
Ga(2)-As(1)-Ga(2')	103.22(7)	102.54(7)
Ga(2)-As(1)-C(111)	111.7(3)	112.1(3)
C(111)-As(1)-C(111')	108.5(4)	108.4(4)
As(1)-Ga(2)-As(3)	121.06(7)	121.65(7)
As(1)-Ga(2)-Br(21)	111.33(7)	112.41(6)
As(1)-Ga(2)-Br(22)	101.76(5)	100.44(5)
As(3)-Ga(2)-Br(21)	111.11(5)	110.92(5)
As(3)-Ga(2)-Br(22)	97.11(6)	96.78(6)
Br(21)-Ga(2)-Br(22)	113.14(8)	112.82(8)
Ga(2)-As(3)-Ga(4)	110.74(5)	110.20(5)
Ga(2)-As(3)-C(311)	98.3(4)	98.4(3)
Ga(2)-As(3)-C(321)	121.7(3)	121.4(3)
Ga(4)-As(3)-C(311)	110.1(3)	110.2(3)
Ga(4)-As(3)-C(321)	107.0(3)	107.8(3)
C(311)-As(3)-C(321)	108.5(5)	108.2(4)
As(3)-Ga(4)-As(3')	113.32(8)	113.59(7)
As(3)-Ga(4)-Br(41)	107.16(4)	109.12(4)
Br(41)-Ga(4)-Br(41')	111.26(9)	111.94(8)
As(1)-C(111)-Si(11)	125.0(4)	125.2(5)
As(3)-C(311)-Si(31)	125.7(7)	125.1(7)
As(3)-C(321)-Si(32)	124.1(6)	123.3(6)

Legends For Figures

Fig. 1. Atomic arrangement in $[(Me_3SiCH_2)_2AsGaBr_2]_3$ (1) (hydrogen atoms omitted for clarity).

Fig. 2. The twist-boat conformation of the $(Ga-As)_3$ ring of $[(Me_3SiCH_2)_2AsGaBr_2]_3$ (1).



C(113')
C(113)
C(114')
C(114)
Si(11)
Si(11')
C(112')
C(112)
C(111)
C(111')
Br(21)
Br(21')
As(1)
Ga(2)
Ga(2')
C(322)
C(322')
C(324)
Si(32)
Si(32')
C(324')
Br(22)
Br(22')
C(323)
C(323')
As(3)
As(3')
C(321)
Ga(4)
C(321')
C(311)
C(311')
C(314)
C(314')
Si(31)
Si(31')
Br(41')
Br(41)
C(312)
C(312')
C(313)
C(313')

Figure 1



Br(22)
C(321')
Br(41')
C(111')
Br(21)
Ga(2)
As(3')
C(311)
As(1)
Ga(4)
Ga(2')
As(3)
C(311')
Br(21')
C(111)
Br(41)
Br(22')
C(321)

Figure 2

DL/1113/87/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1
Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

END
DATE
9-88
DTIC